

CANONICAL COMMUTATION RELATIONS IN THE SCHWINGER MODEL¹

Gil Gat and O.W. Greenberg²

*Center for Theoretical Physics
Department of Physics and Astronomy
University of Maryland
College Park, MD 20742-4111*

Preprint number 94-087

Abstract

We give the first operator solution of the Schwinger model that obeys the canonical commutation relations in a covariant gauge.

1. Introduction

The Schwinger model[1, 2], massless two-dimensional quantum electrodynamics, is an exactly solvable model that exhibits confinement analogous to the confinement that is thought to occur in four-dimensional quantum chromodynamics (QCD). Because of this, the model has been extensively discussed in the literature using both operator and path integral techniques. Surprisingly, the published operator solutions do not obey the canonical commutation relations that follow from the Lagrangian. The purpose of this letter is to provide an operator solution that does obey these relations.

2. Review of the Lowenstein-Swieca solution of the massless Schwinger model

The Schwinger model is quantum electrodynamics in 1+1 dimensions. In the

¹Supported in part by the National Science Foundation, e-mail addresses: gat@wam.umd.edu, greenberg@umdep1.umd.edu

²Supported in part by a Semester Research Grant from the University of Maryland, College Park.

Lorentz gauge,³

$$\mathcal{L} = \bar{\psi} i \not{D} \psi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{2} (\partial \cdot A)^2, \quad (1)$$

where

$$D^\mu = \partial^\mu - e A^\mu, \quad F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu. \quad (2)$$

In its massless form this model is exactly solvable. Lowenstein and Swieca[3] found an operator ansatz that yields the matrix elements computed by Schwinger. Their ansatz solution has the following form:

$$A^\mu(x) = -\sqrt{\frac{\pi}{e^2}} (\epsilon^{\mu\nu} \partial_\nu \Sigma + \partial^\mu \eta), \quad (3)$$

$$\psi(x) =: \exp[i\sqrt{\pi}(\gamma_5 \Sigma(x) - \eta(x))] : \psi^{(0)}(x), \quad (4)$$

where η is a free neutral massless field with negative metric corresponding to the gauge degrees of freedom satisfying $[\eta(x), \dot{\eta}(y)]_{ET} = -i\delta(x^1 - y^1)$, while Σ is a free neutral massive field with positive metric representing the physical degrees of freedom satisfying $[\Sigma(x), \dot{\Sigma}(y)]_{ET} = i\delta(x^1 - y^1)$ and $\psi^{(0)}$ is a solution of the free massless Dirac equation. (ET stands for equal time.) This solution displays the main property of the Schwinger model: the only physical state is a free particle of mass $e/\sqrt{\pi}$. Lowenstein and Swieca did not discuss the canonical commutation relations since their interest was to get Schwinger's matrix elements.

Capri and Ferrari[4] found another solution of the model, in covariant gauges; however, this solution does not obey the canonical commutation relations. Both solutions use ansatzes to solve the operator equations; while this is satisfactory, we believe it can be illuminating to solve the equations in a more direct way, and a direct method of solution may be helpful in analyzing the massive model and other models. The purpose of this letter is to provide a direct solution that obeys the canonical commutation relations. We recognize that this is a rather technical matter; however, in view of the extensive literature on this model, we think the literature should contain a solution that obeys these relations.

³This form of the Lagrangian ignores wavefunction renormalization of the spinor field. Taking account of this wavefunction renormalization, the spinor term in the Lagrangian is $\bar{\psi}_u i \not{D} \psi_u - \langle \bar{\psi}_u i \not{D} \psi_u \rangle_0 = Z \bar{\psi} i \not{D} \psi$, where ψ_u is the unrenormalized spinor field and Z is the spinor wavefunction renormalization. We suppress the factor Z below.

3. Direct operator solution of the Schwinger model

We use the Lagrangian Eq.(1), but drop surface terms so that the gauge field part becomes

$$-\frac{1}{2}(\partial_\mu A_\nu)(\partial^\mu A^\nu). \quad (5)$$

Since the Lorentz group (without inversions) is abelian in 1+1, all irreducible representations are one-dimensional; thus the vector and spinor fields in the model are composed of one-dimensional irreducibles arbitrarily pasted together. We choose to express the Lagrangian in terms of the irreducible fields in the basis in which

$$A^0 = \frac{1}{2}(A^+ + A^-), \quad A^1 = \frac{1}{2}(A^+ - A^-), \quad \psi = (\psi_1, \psi_2), \quad (6)$$

with

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^1 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (7)$$

In terms of the irreducible fields,

$$\mathcal{L} = \psi_1^\dagger(2i\partial^+ - eA^-)\psi_1 + \psi_2^\dagger(2i\partial^- - eA^+)\psi_2 + \frac{1}{2}(\partial^1 A^+ \partial^1 A^- - \partial^0 A^+ \partial^0 A^-). \quad (8)$$

where we have introduced lightcone coordinates, $x^+ = x^0 + x^1$, $x^- = x^0 - x^1$. The corresponding derivatives are $\frac{\partial}{\partial x^\pm} = \frac{1}{2}(\frac{\partial}{\partial x^0} \pm \frac{\partial}{\partial x^1})$. We define these so that $\frac{\partial}{\partial x^\pm} x^\pm = 1$. Note that although the fields $A^\pm$ are lightcone fields, we are *not* using lightcone quantization, but rather are using equal-time canonical quantization. The naive operator equations of motion are

$$\square A^+ - 2e\psi_1^\dagger\psi_1 = 0, \quad (9)$$

$$\square A^- - 2e\psi_2^\dagger\psi_2 = 0, \quad (10)$$

$$(2i\partial^+ - eA^-)\psi_1 = 0, \quad (11)$$

$$(2i\partial^- - eA^+)\psi_2 = 0. \quad (12)$$

As Schwinger pointed out in his original paper, the spinor bilinear products require a line integral of the “vector” potential between the $\psi^\dagger$ and the ψ in order to ensure gauge invariance; this is done explicitly below using point-splitting. For example,

$\psi_1^\dagger \psi_1$ is replaced by

$$\lim_{\epsilon \rightarrow 0} \frac{1}{2} \left[\psi_1^\dagger(x + \epsilon) e^{-ie \int_x^{x+\epsilon} A_\mu(w) dw^\mu} \psi_1(x) + cc. \right]. \quad (13)$$

The canonical momenta are

$$\pi_{A^+} = -\frac{1}{2} \partial^0 A^-, \quad \pi_{A^-} = -\frac{1}{2} \partial^0 A^+, \quad \pi_{\psi_j} = i\psi_j^\dagger. \quad (14)$$

First we solve the Dirac equations by exponentiation,

$$\psi_1(x) = \mathcal{P} \exp \left[-\frac{ie}{2} \int_{-\infty}^{x^+} A^-(w^+, x^-) dw^+ \right] \psi_1^{(0)}(x^-), \quad \partial^+ \psi_1^{(0)} = 0, \quad (15)$$

$$\psi_2(x) = \mathcal{P} \exp \left[-\frac{ie}{2} \int_{-\infty}^{x^-} A^+(x^+, w^-) dw^- \right] \psi_2^{(0)}(x^+), \quad \partial^- \psi_2^{(0)} = 0. \quad (16)$$

The point-splitting vector is taken spacelike, $\epsilon = (0, \epsilon^1)$, $\epsilon^\pm = \pm \epsilon^1$. Thus, for example, $\psi_1^\dagger$ must be replaced by

$$\psi_1^\dagger(x + \epsilon) = \psi_1^{(0)\dagger}(x^- - \epsilon^1) \bar{\mathcal{P}} \exp \left[\frac{ie}{2} \int_{-\infty}^{x^+ + \epsilon^1} A^-(w^+, x^- - \epsilon^1) dw^+ \right]. \quad (17)$$

The symbols $\mathcal{P}$ and $\bar{\mathcal{P}}$ stand for path and antipath ordering, respectively. The result of the point-splitting differs from the usual one by having integrated (nonlocal) terms. The equations for $A^\pm$ become

$$(\square + \frac{e^2}{2\pi}) A^+ - \frac{e^2}{2\pi} \int_{-\infty}^{x^+} \frac{\partial A^-}{\partial x^-}(w^+, x^-) dw^+ = 2e\psi_1^{(0)\dagger}(x^-) \psi_1^{(0)}(x^-) \quad (18)$$

$$(\square + \frac{e^2}{2\pi}) A^- - \frac{e^2}{2\pi} \int_{-\infty}^{x^-} \frac{\partial A^+}{\partial x^+}(x^+, w^-) dw^- = 2e\psi_2^{(0)\dagger}(x^+) \psi_2^{(0)}(x^+). \quad (19)$$

The integrated terms here can be removed by taking derivatives with respect to the upper limit. Combining the resulting equations leads to

$$\square \cdot A = 0 \quad (20)$$

$$(\square + \frac{e^2}{\pi}) \epsilon_{\mu\nu} \partial^\mu A^\nu = 0; \quad (21)$$

thus $\partial \cdot A \equiv \eta$ is a massless field and $\epsilon_{\mu\nu}\partial^\mu A^\nu \equiv \Sigma$ is a field of mass $e/\sqrt{\pi}$. As Lowenstein and Swieca observed, η and Σ (which is the electric field in 1+1) are the gauge-variant and gauge-invariant degrees of freedom, respectively. Then $\square A^\mu$ must be a linear combination of $\partial^\mu \eta$ and $\epsilon^{\mu\nu}\partial_\nu \Sigma$. A^μ must be the convolution of the $\bar{\Delta}(x) = -\frac{1}{2}\epsilon(x^0)\Delta(x)$ Green's function with this linear combination plus terms annihilated by $\square$. The convolution of $\bar{\Delta}(x)$ with η does not exist, because, formally, it is $\int d^2y \bar{\Delta}(x-y)\eta(y) = \int d^2k \exp(-ik \cdot x) \delta(k^2) \tilde{\eta}(k)/k^2$, which is ill-defined. Because of this, a new field a that obeys $\square a = \eta$ must be introduced. This was first done by Capri and Ferrari. Thus

$$A^\mu = c_1 \partial^\mu a + c_2 \epsilon^{\mu\nu} \partial_\nu \Sigma + c_3 \partial^\mu \eta + c_4 \bar{\psi}^{(0)} \gamma^\mu \psi^{(0)}. \quad (22)$$

For the massless case,

$$\bar{\psi}^{(0)} \gamma^\mu \psi^{(0)} = \partial^\mu \phi, \quad \square \phi = 0, \quad (23)$$

where ϕ is a free positive-metric scalar field. Substitution of this A^μ into the integrodifferential Eqs.(18,19) fixes $c_4 = -2e/\pi$ and leaves the other constants arbitrary.

We want the equal-time canonical commutation relations (for A^μ they are not renormalized), that play the role of boundary conditions, to determine these constants. To do this, we must find the equal-time relations among a and η and their canonical conjugates. In order to determine these relations, we calculate the two-point functions of the fields η and a and assume they are asymptotic fields with c-number commutators. The coefficients c_i , $i = 1, 2, 3$ in A^μ take care of the absolute normalizations; we allow a metric factor $\tau = \pm 1$ in the two-point function of η and let the CCR's determine the metrics of η and a . We use Klaiber's regulation[5] of the massless scalar two-point function,

$$\langle 0 | \eta(x) \eta(y) | \rangle = \frac{\tau}{2\pi} \int_{-\infty}^{\infty} \frac{dp^1}{2|p^1|} [e^{-ip \cdot \xi} - \theta(\lambda - |p^1|)] \quad (24)$$

$$= -\frac{\tau}{4\pi} \ln(-\mu^2 \xi^2 + i\xi^0 \epsilon) \quad (25)$$

$$\equiv D^{(+)}(\xi; \lambda), \quad (26)$$

where $\xi = x - y$, $\mu = e^\gamma \lambda$, $\gamma = \text{Euler's const.}$ From $\square \langle 0 | a(x) \eta(y) | 0 \rangle = \langle 0 | \eta(x) \eta(y) | 0 \rangle$, we find

$$\langle 0 | a(x) \eta(y) | \rangle = \tau I^{(+)}(\xi; \lambda) + c_{a\eta} D^{(+)}(\xi; \lambda), \quad (27)$$

$$I^{(+)}(\xi; \lambda) = -\frac{1}{4\pi} \left[\frac{\xi^2}{4} \ln(-\mu^2 \xi^2 + i\epsilon \xi^0) - \frac{1}{2} \xi^2 \right], \quad (28)$$

where the $D^{(+)}$ term is an arbitrary solution of the homogeneous equation. (Capri and Ferrari's result for $I^{(+)}$ is incorrect). Analogously,

$$\langle 0 | \eta(x) a(y) | \rangle = \tau I^{(+)}(\xi; \lambda) + c_{\eta a} D^{(+)}(\xi; \lambda). \quad (29)$$

The two-point function of two a 's can be calculated in two ways; consistency requires $c_{a\eta} = c_{\eta a}$. Then

$$\langle 0 | a(x) a(y) | \rangle = \tau K^{(+)}(\xi) + c_{a\eta} I^{(+)}(\xi; \lambda) + c_{aa} D^{(+)}(\xi; \lambda), \quad (30)$$

$$K^{(+)}(\xi) = -\frac{1}{256\pi} (\xi^2)^2 [\ln(-\mu^2 \xi^2 + i\epsilon \xi^0) - 3], \quad (31)$$

where we introduce a new function, $K^{(+)}$. The ETCR's follow simply from the assumption that the a and η fields are free.

$$[\eta, \dot{\eta}]_{ET} = \tau i\delta, \quad [\eta, \dot{a}]_{ET} = c_{a\eta} i\delta, \quad [a, \dot{\eta}]_{ET} = c_{a\eta} i\delta, \quad [a, \dot{a}]_{ET} = c_{aa} i\delta, \quad (32)$$

The extra terms that arise from solutions of the homogeneous equations for the two-point functions allow a lot of freedom. The system is much more constrained if these terms are all set to zero, i.e., $c_{a\eta} = c_{aa} = 0$; in this case there are two solutions that obey the CCR's:

$$\tau = -1, \quad c_1 = \mp \sqrt{\frac{e^2}{5\pi}}, \quad c_2 = \pm \sqrt{\frac{\pi}{e^2}}, \quad c_3 = \pm \sqrt{\frac{5\pi}{e^2}}. \quad (33)$$

The sign ambiguity is due to the fact that the CCR's are bilinear in A^μ and $\dot{A}^\mu$. As expected, $\tau = -1$ shows that the fields η and a that are associated with the gauge-variant degree of freedom are ghosts.

4. Conclusions

The canonical commutation relations in field theory and their predecessors in classical mechanics and quantum mechanics are important for many reasons. The Poisson brackets in classical mechanics, for example, ensure that the Hamiltonian is the generator of time translations. In quantum mechanics, for example, the relation $[x, p] = i\hbar$ leads to the uncertainty relation. In quantum field theory, the CCR's lead

to the free field being a collection of quantized oscillators. In nonrelativistic field theories at least, the CCR's imply unitarity[6]. A new feature of the canonical commutation relations in quantum field theory is that they ensure that the asymptotic fields have the proper free commutation relation. (The renormalized canonical commutation relations will do as well as the original CCR's for this purpose.) For these reasons, a solution that obeys the canonical commutation relations is important.

Acknowledgements

It is a pleasure to thank Joseph Sucher for reading the manuscript and for making good suggestions.

References

- [1] J. Schwinger in *Theoretical Physics*, ed. A. Salam (International Atomic Energy Agency, Vienna, 1963), p89. and *Phys. Rev* 128 (1962) 2425.
- [2] *Two Dimensional Quantum Field Theory*, E. Abdalla, M.C. Abdalla and K.D. Rothe (World Scientific, Singapore, 1991) contains an extensive list of references on the Schwinger model.
- [3] J. Lowenstein and J. A. Swieca *Ann. of Phys.* 68 (1971) 172.
- [4] A.Z Capri and R. Ferrari, *Nuovo Cimento* 62A (1981) 273 and *J. Math. Phys.* 25 (1984) 141.
- [5] B. Klaiber, in *Lectures in Theoretical Physics, Vol X-A, Quantum Theory and Statistical Physics*, ed. A.O. Barut and W.E. Brittin (Gordon and Breach, New York, 1968), p141.
- [6] S. Corley and O.W. Greenberg, University of Maryland Preprint No. 94-094